

# TREE PEST UPDATES

75 Santa Barbara Rd, 2nd floor, Pleasant Hill, CA 94523-4215 (925) 646-6540  
 Diablo Valley Farm Center, Delta Rd. & 2<sup>nd</sup> St., Knightsen (no phone yet!)

April 12, 2005

## CODLING MOTH

**HOST CROPS:** Apple, Pear, Walnut

*[Not all walnut orchards need treatment every year or every generation. If you had greater than 3% worm damage last year and/or are catching high numbers in your traps, you probably need to treat this generation.]*

**BIOFIX 1A:** I am establishing our areawide biofix as **March 30<sup>th</sup>** – this is the first day that we had flight in most orchards and consistent weather suitable for mating to occur. *Use the date that the flight began in your own traps and sunset temperatures were at least 62F as your own orchard biofix.*

**TREATMENT OPTIONS:** If the weather remains “normal,” the projected treatment dates are:

*Organophosphate (OP) Sprays (Guthion, Imidan, Lorsban, Penncap, Supracide):*

Apples should be sprayed at 250 degree days (DD). This year this should occur about **April 26<sup>th</sup> OR 28 calendar days after your own orchard biofix**. You may need to reapply in 14 days if flights are extended and population pressure is high in your orchard.

Walnuts have more lenient damage standards and can wait until 300-400 DD. This should occur about **April 30<sup>th</sup>-May 7<sup>th</sup> OR 32-39 days after your own orchard biofix**. If you have low pressure you may want to wait until the 1B flight (late May) or 2A flight (late June) to apply a spray on walnuts, if needed.

*“Reduced Risk” Materials (Assail, Intrepid, Success, Last Call):* These materials are soft on beneficials and less toxic to people and the environment. Assail and Intrepid are the newer and more effective materials. Intrepid and Success should be applied at the beginning of egg hatch about 200 DD. This should occur about **April 21<sup>st</sup> OR 23 days after your own biofix**. Assail should be applied at the traditional OP timing (above). All these materials may need reapplication after 10-18 days if flights are extended and population pressure is high. Last Call should be applied at or shortly after biofix and reapplied at 4-5 week intervals.

*Mating Disruption:* Dispensers should have been hung in the orchards **before March 30<sup>th</sup>** OR your own orchard biofix. If they went up afterwards, you may want to consider a supplemental control for the first generation.

*Organic Sprays (Entrust, Surround, Oil):* Entrust is the new organic formulation of Success. It is the most effective organic spray currently available. It should be applied at 200 DD (**April 21<sup>st</sup> OR 23 days after your own biofix**) and reapplied at 10 day intervals, if continued coverage is needed. Oil and Surround are mild controls and best used as a supplement to Mating Disruption. They should be re-applied every 7-14 days in 100-200 gallons of water/acre. Begin Surround applications at 100 DD (**April 11 OR 12 days after your own biofix**) to discourage egg laying. Begin Oil applications at 200 DD (**April 21<sup>st</sup> OR 23 days after your own biofix**) to suffocate eggs. *Do not apply oil within a few weeks of a sulfur application.*

Note: The above information is provided to serve as baseline data for East Contra Costa County. For best results compare with traps and observations in your own orchards. Depending on pest pressure, sprays may not be necessary. Projected treatment times are based on historical weather data.

## **CODLING MOTH UPDATE**

UC Cooperative Extension  
75 Santa Barbara Rd, 2nd floor  
Pleasant Hill, CA 94523

**FOR MORE DETAILED INFORMATION ON TREATMENTS AND TIMING, CALL FOR A COPY OF OUR MOST RECENT CODLING MOTH IPM GUIDELINES FOR APPLES, PEARS, OR WALNUTS.**

*Many thanks to Suterra for providing BioLure traps for the Contra Costa County Tree Pest Update Program.*

Janet Caprile  
Farm Advisor

The University of California prohibits discrimination or harassment of any person on the basis of race, color, national origin, religion, sex, gender identity, pregnancy (including childbirth, and medical conditions related to pregnancy or childbirth), physical or mental disability, medical condition (cancer-related or genetic characteristics), ancestry, marital status, age, sexual orientation, citizenship, or status as a covered veteran (covered veterans are special disabled veterans, recently separated veterans, Vietnam era veterans, or any other veterans who served on active duty during a war or in a campaign or expedition for which a campaign badge has been authorized) in any of its programs or activities. University policy is intended to be consistent with the provisions of applicable State and Federal laws. Inquiries regarding the University's nondiscrimination policies may be directed to the Affirmative Action/Staff Personnel Services Director, University of California, Agriculture and Natural Resources, 300 Lakeside Drive, 6<sup>th</sup> Floor, Oakland, CA 94612-3550, (510) 987-0096.

**What's a Biofix?:** It's just the beginning of the flight for each new generation. We usually have 3 generations for codling moth in this area. We use the Biofix to begin degree day calculations for each generation so we know when egg laying, hatchout, and other lifecycle events will happen. This helps us to time our treatments most effectively.

**What's a Degree Day?** Insects develop faster or slower depending on the temperature. Degree days are a measure based on the maximum and minimum temperatures for each day which allow us to figure out how fast the insects are developing. You may see them abbreviated as DD or °D. If you have the daily maximum & minimum temperatures for your orchard, you can look the degree days up on a chart. If you have access to the Internet, you can get Brentwood weather data and do a degree day calculation from the UC IPM Program home page. This page also lets you calculate the projected degree days based on historical weather data so you can make projections for treatment windows (this is how I do it!). The address is <http://www.ipm.ucdavis.edu>. Give me a call if you would like a